

著作目録（平則夫）

著者	東北大学史料館
号	563
発行年	1995-03
URL	http://hdl.handle.net/10097/00065380

平 則 夫 教 授 著 作 目 録

平 成 7 年 3 月
東 北 大 学 記 念 資 料 室
(著 作 目 録 第 563 号)



平 則 夫 教 授 略 歴

生年月日 昭和6年11月14日生
出生地 宮城県
所 属 東北大学医学部薬理学第二講座

学 歴

昭和31年3月 東北大学医学部卒業
昭和36年3月 東北大学大学院医学研究科生理学系専攻博士課程修了

学 位

昭和36年3月 医学博士（東北大学）

受 賞

な し

職 歴

昭和36年4月 東北大学医学部助手
昭和38年4月 東北大学医学部講師
昭和39年2月 東北大学医学部助教授
昭和39年9月 China Medical Board Fellow として薬理学に関する研究のためアメリカ合衆国に出張（昭和41年3月まで）
昭和47年1月 東北大学医学部教授
平成3年4月 東北大学医学部学部長
平成7年3月 定年退職

学会活動

日本生理学会評議員	昭和38年4月～
日本薬理学会評議員	昭和44年4月～
日本薬理学会理事	昭和53年5月～昭和57年4月，昭和61年5月～平成2年4月， 平成4年4月～平成5年3月
日本薬理学会副会長	平成2年4月～平成3年3月
日本薬理学会会長	平成3年4月～平成4年3月
日本臨床薬理学会評議員	昭和55年2月～
日本循環器学会評議員	昭和57年4月～
日本心電学会運営委員	昭和59年1月～
日本麻酔・薬理学会理事	昭和63年6月～
日本循環制御学会評議員	平成3年12月～

著 作 目 録

Original papers

- 1) Suzuki, H. and *Taira, N.*: Regional difference of the direct cortical response. Jpn. J. Physiol. 8: 365-377, 1958.
- 2) Suzuki, H. and *Taira, N.*: An analysis of direct cortical response. Tohoku J. Exp. Med. 70: 1-10, 1959.
- 3) Suzuki, H., *Taira, N.* and Motokawa, K.: Spectral response curves and receptive fields of pre- and postgeniculate fibres of the cat. Tohoku J. Exp. Med. 71: 401-415, 1960.
- 4) *Taira, N.*: Origin of hippocampal seizure. Tohoku J. Exp. Med. 73: 191-199, 1961.
- 5) Suzuki, H. and *Taira, N.*: Effect of reticular stimulation upon synaptic transmission in cat's lateral geniculate body. Jpn. J. Physiol. 11: 641-655, 1961.
- 6) *Taira, N.* and Okuda, J.: Sensory transmission in visual pathway in various arousal states of cat. Tohoku J. Exp. Med. 78: 76-96, 1962.
- 7) Okuda, J., *Taira, N.* and Motokawa, K.: Spectral response curves of postgeniculate neurons in the cat. Tohoku J. Exp. Med. 78: 147-157, 1962.
- 8) Motokawa, K., *Taira, N.* and Okuda, J.: Spectral responses of single units in the primate visual cortex. Tohoku J. Exp. Med. 78: 320-337, 1962.
- 9) Hashimoto, K., Kumakura, S. and *Taira, N.*: Vascular reflex responses induced by an intraarterial injection of azaazepinophenothiazine, andromedotoxin, veratridine, bradykinin and kallikrein and blocking action of sodium salicylate. Jpn. J. Physiol. 14: 299-308, 1964.
- 10) *Taira, N.*, Imazawa, Y. and Motokawa, K.: Electrical stimulation of on- and off-units in cat's retina. Tohoku J. Exp. Med. 85: 89-104, 1965.
- 11) Tominaga, S., Miura, M., *Taira, N.* and Hashimoto, K.: Reactive hyperemia of hindlimb after the treatment with IAc, FAc and DNP. Jpn. J. Physiol. 16: 1-10, 1966.
- 12) Hirata, M., Tanaka, S., *Taira, N.* and Hashimoto, K.: Comparative studies on pseudodffective responses to different algesic substances in conscious dogs. The effect of sodium salicylate. Arch. Int. Pharmacodyn. Ther. 162: 150-161, 1966.

- 13) *Taira, N.* and Marshall, J. M.: Effects of methylguanidine on the electrical activity of single pacemaker fibers in the sinus node of the rabbit. *J. Pharmacol. Exp. Ther.* 155: 69-75, 1967.
- 14) *Taira, N.* and Marshall, J. M.: Action of oxytocin antagonists on electrical and mechanical activity of the uterus. *Am. J. Physiol.* 212: 725-731, 1967.
- 15) *Taira, N.*, Nakayama, K., Sano, N. and Hashimoto, K.: Nicotinic property of acetylcholine receptors subserving nociception in the puppy hindlimb as revealed by the activity of cholinergic agonists administered into the femoral artery. *Tohoku J. Exp. Med.* 96: 111-112, 1968.
- 16) Yasuda, K., Satoh, S., *Taira, N.* and Hashimoto, K.: Potentiation of norepinephrine and adenosine in the renal circulation by 2,6-bis (diethanolamino) -4,8-dipiperidino-pyrimido (5,4-d) pyrimidine (dipyridamole) treatment (Preliminary report). *Tohoku J. Exp. Med.* 96: 161-164, 1968.
- 17) *Taira, N.*, Matsumura, S. and Hashimoto, K.: Excitation of the parasympathetic ganglia of the canine urinary bladder by a muscarinic ganglionic stimulant, 4-(*m*-chlorophenyl-carbamoyloxy)-2-butyryl-trimethylammonium chloride (McN-A-343). *Tohoku J. Exp. Med.* 96: 209-210, 1968.
- 18) Matsumura, S., *Taira, N.* and Hashimoto, K.: The pharmacological behavior of the urinary bladder and its vasculature of the dog. *Tohoku J. Exp. Med.* 96: 247-258, 1968.
- 19) *Taira, N.*, Nakayama, K. and Hashimoto, K.: Vocalization response of puppies to intra-arterial administration of bradykinin and other algescic agents, and mode of actions of blocking agents. *Tohoku J. Exp. Med.* 96: 365-377, 1968.
- 20) *Taira, N.*, Yamamoto, M., Satoh, S. and Hashimoto, K.: Absence of blocking effect of tetrodotoxin on the action of tyramine in the saphenous vascular bed of the dog. *Tohoku J. Exp. Med.* 97: 97-98, 1969.
- 21) *Taira, N.*, Matsumura, S. and Hashimoto, K.: Effect of tetrodotoxin on the bladder response to pelvic nerve stimulation and intra-arterial 1,1-dimethyl-4-phenylpiperazinium and acetylcholine in the dog. *Tohoku J. Exp. Med.* 97: 283-288, 1969.
- 22) Sakai, K., Yasuda, K., *Taira, N.* and Hashimoto, K.: α -Adrenergic blocking action of norephedrine-theophylline (NET) observed in the renal vascular response. *Jpn. J. Pharmacol.* 19: 194-198, 1969.

- 23) Hashimoto, K., Matsumura, S., Sano, N. and *Taira, N.*: Comparison of β -adrenergic blocking activity of dichloroisoprenaline, H 56/28, I.C.I. 50172, LB 46, methoxamine, MJ 1999 and propranolol in the canine femoral vascular bed. *Experientia* 25: 1155-1156, 1969.
- 24) Hashimoto, K., Ohkuda, K., Chiba, S. and *Taira, N.*: Beta-adrenergic blocking effect of dichloroisoprenaline (DCI), H 56/28, I.C.I. 50172, LB 46, methoxamine, MJ 1999 and propranolol on the sinus node activity of the dog heart. *Experientia* 25: 1156-1157, 1969.
- 25) Yamamoto, M., *Taira, N.* and Hashimoto, K.: Cholinergic mechanism for intestinal hypermotility caused by arterial occlusion. *Jpn. J. Physiol.* 20: 145-153, 1970.
- 26) Ohkuda, K., Chiba, S., *Taira, N.* and Hashimoto, K.: Comparison of beta-adrenergic blocking effect of DCI, methoxamine, propranolol, MJ 1999, H 56/28, LB 46 and I.C.I. 50172 on the chronotropic response to isoprenaline. *Jpn. J. Pharmacol.* 20: 294-304, 1970.
- 27) Hashimoto, K., Endoh, M., Tamura, K. and *Taira, N.*: Comparison of β -adrenergic blocking activity of DCI, H56/28, ICI50172, LB46, methoxamine, MJ1999 and propranolol in the blood perfused canine papillary muscle preparation. *Experientia* 26: 757-759, 1970.
- 28) Yamamoto, M., Sasaki, H., *Taira, N.* and Hashimoto, K.: Responses of the canine saphenous vascular bed to various biogenic and foreign substances. *Tohoku J. Exp. Med.* 101: 299-309, 1970.
- 29) *Taira, N.*, Nakayama, K. and Hashimoto, K.: Nicotinic acetylcholine receptors subserving nociception in the dog hindlimb. *Jpn. J. Physiol.* 20: 571-583, 1970.
- 30) *Taira, N.*, Matsumura, S. and Hashimoto, K.: Excitation of the parasympathetic ganglia of the canine urinary bladder through a muscarinic mechanism. *J. Pharmacol. Exp. Ther.* 176: 93-100, 1971.
- 31) Hashimoto, K., *Taira, N.*, Hirata, M. and Kokubun, M.: The mode of hypotensive action of newly synthesized sydnonimine derivatives. *Arzneim.-Forsch.* 21: 1329-1332, 1971.
- 32) Hashimoto, K., Iijima, T., *Taira, N.* and Hashimoto, K.: Negative dromotropic response to phenylephrine blocked by phenoxybenzamine. *Tohoku J. Exp. Med.* 105: 305-307, 1971.

- 33) *Taira, N.*, Iijima, T., Hashimoto, K. and Hashimoto, K.: A device for automatic plotting of the atrioventricular conduction time. *Tohoku J. Exp. Med.* 105: 391-395, 1971.
- 34) Hashimoto, K., *Taira, N.*, Chiba, S., Hashimoto, K., Jr., Endoh, M., Kokubun, M., Kokubun, H., Iijima, T., Kimura, T., Kubota, K. and Oguro, K.: Cardiohemodynamic effects of BAY a 1040 in the dog. *Arzneim.-Forsch.* 22: 15-21, 1972.
- 35) Nakayama, K., *Taira, N.* and Hashimoto, K.: Absence of relation between the vocalization response and the intestinal motility induced by intra-arterial administration of aza-azepinophenothiazine (RPP-201) into the mesenteric artery of the dog. *Tohoku J. Exp. Med.* 106: 271-274, 1972.
- 36) Nakayama, K., *Taira, N.*, Hashimoto, K., Kobayashi, S., Kumakura, S., Oshima, T., Koike, H. and Takagi, H.: Effects of anisotropine methylbromide (Valpin®) and its mixture with sulpyrine on vocalization response and spasm of intestine induced by acetylcholine in dogs. *Jpn. J. Pharmacol.* 22: 215-220, 1972.
- 37) Iijima, T., Motomura, S., *Taira, N.* and Hashimoto, K.: Effects on A-V conduction of tetrodotoxin selectively injected into the A-V node artery of the dog. *Tohoku J. Exp. Med.* 107: 99-100, 1972.
- 38) *Taira, N.*, Iwatsuki, K., Hashimoto, K. and Hirayama, H.: Effect of ivory shell toxin on the parasympathetic ganglia of the dog urinary bladder. *Tohoku J. Exp. Med.* 107: 191-192, 1972.
- 39) Hashimoto, K., Iijima, T., Hashimoto, K. and *Taira, N.*: The isolated and cross-circulated AV node preparation of the dog. *Tohoku J. Exp. Med.* 107: 263-275, 1972.
- 40) Kimura, T., Endoh, M., *Taira, N.* and Hashimoto, K.: Comparison of new β -adrenergic blockers, C-3, K61366, Y-6124 and YB-2, with pindolol and propranolol in the blood perfused canine papillary muscle preparation. *Experientia* 28: 813-814, 1972.
- 41) Hashimoto, K., Kubota, K., Chiba, S. and *Taira, N.*: Comparison of β -adrenergic blocking activity of eight blockers in the excised and blood-perfused canine sino-atrial node preparation. *Experientia* 28: 822-823, 1972.
- 42) *Taira, N.* and Hashimoto, K.: Absence of the pain-producing activity of otherwise biologically active synthetic fragments of bradykinin. *Tohoku J. Exp. Med.* 110: 191-194, 1973.

- 43) Iijima, T., Hashimoto, K., *Taira, N.* and Hashimoto, K.: Dromotropic effects of optical isomers of *beta* blockers on the cross-circulated canine atrioventricular node preparation. *J. Pharmacol. Exp. Ther.* 186: 145-151, 1973.
- 44) Satoh, S., *Taira, N.* and Hashimoto, K.: Absence of a neurally-mediated process in salivation and vasodilation caused by McN-A-343 in the dog submaxillary gland. *Jpn. J. Pharmacol.* 23: 581-583, 1973.
- 45) *Taira, N.* and Satoh, S.: Prostaglandin $F_{2\alpha}$ as a potent excitant of the parasympathetic postganglionic neurons of the dog salivary gland. *Life Sci.* 13: 501-506, 1973.
- 46) *Taira, N.* and Nakano, T.: Is intra-arterial 5-hydroxytryptamine a potent algogenic substance in the dog? *Eur. J. Pharmacol.* 25: 113-116, 1974.
- 47) Kokubun, M., *Taira, N.* and Hashimoto, K.: Cardiohemodynamic effects of nitroglycerin and several vasodilators. *Jpn. Heart J.* 15: 126-144, 1974.
- 48) Oguro, K., *Taira, N.* and Hashimoto, K.: Cardiohemodynamic effect of 5-methyl-7-diethylamino-s-triazolo-(1,5-a) pyrimidine (Trapymine). *Arzneim.-Forsch.* 24: 911-914, 1974.
- 49) Iijima, T., Motomura, S., *Taira, N.* and Hashimoto, K.: Selective suppression of neural excitation by tetrodotoxin injected into the canine atrioventricular node artery. *J. Pharmacol. Exp. Ther.* 189: 638-645, 1974.
- 50) Iijima, T., Motomura, S., *Taira, N.* and Hashimoto, K.: Comparison of effects of glucagon and isoprenaline on atrio-ventricular conduction and sino-atrial rate in the dog heart. *Clin. Exp. Pharmacol. Physiol.* 1: 241-248, 1974.
- 51) *Taira, N.* and Satoh, S.: Differential effects of tetrodotoxin on the sialogenous and vasodilator actions of prostaglandin E_2 in the dog salivary gland. *Life Sci.* 15: 987-993, 1974.
- 52) Sakai, K., Sugano, S., *Taira, N.* and Hashimoto, K.: Pharmacological features of peripheral vascular beds of beagles. *Jpn. J. Pharmacol.* 24: 659-669, 1974.
- 53) *Taira, N.*: Mode of actions of prostaglandin $F_{2\alpha}$ on the urinary bladder and its arterial bed in the dog. *Eur. J. Pharmacol.* 29: 30-34, 1974.
- 54) Himori, N., Ono, H. and *Taira, N.*: Dual effects of a new coronary vasodilator, diltiazem, on the contractile force of the blood-perfused papillary muscle of the dog. *Jpn. J. Pharmacol.* 25: 350-352, 1975.

- 55) Motomura, S., Iijima, T., *Taira, N.* and Hashimoto, K.: Effects of neurotransmitters injected into the posterior and the anterior septal artery on the automaticity of the atrioventricular junctional area of the dog heart. *Circ. Res.* 37: 146-155, 1975.
- 56) *Taira, N.* and Narimatsu, A.: Effects of nifedipine, a potent calcium-antagonistic coronary vasodilator, on atrioventricular conduction and blood flow in the isolated atrioventricular node preparation of the dog heart. *Naunyn-Schmiedeberg's Arch. Pharmacol.* 290: 107-112, 1975.
- 57) *Taira, N.*, Narimatsu, A. and Satoh, S.: Differential block by *l*-hyoscyamine of the salivary and vascular responses of the dog mandibular gland to prostaglandin F_{2a} . *Life Sci.* 17: 1869-1875, 1975.
- 58) *Taira, N.*, Motomura, S., Narimatsu, A. and Iijima, T.: Experimental pharmacological investigations of effects of nifedipine on atrioventricular conduction in comparison with those of other coronary vasodilators. In: 2nd International Adalat® Symposium — New Therapy of Ischemic Heart Disease, ed. by W. Lochner, W. Braasch and G. Kroneberg, pp. 40-48, Springer — Verlag, Berlin — Heidelberg — New York, 1975.
- 59) Himori, N. and *Taira, N.*: Effects of quinidine on blood flow rate and developed tension in blood-perfused canine papillary muscle. *Clin. Exp. Pharmacol. Physiol.* 3: 1-7, 1976.
- 60) Himori, N. and *Taira, N.*: A method for recording smooth muscle and vascular responses of the blood-perfused dog trachea *in situ*. *Br. J. Pharmacol.* 56: 293-299, 1976.
- 61) Iijima, T. and *Taira, N.*: Modification by manganese ions and verapamil of the responses of the atrioventricular node to norepinephrine. *Eur. J. Pharmacol.* 37: 55-62, 1976.
- 62) Nakano, T. and *Taira, N.*: 5-Hydroxytryptamine as a sensitizer of somatic nociceptors for pain-producing substances. *Eur. J. Pharmacol.* 38: 23-29, 1976.
- 63) Himori, N. and *Taira, N.*: Release of noradrenaline proposed as a mechanism for the positive inotropic action of dipyridamole. *Naunyn-Schmiedeberg's Arch. Pharmacol.* 294: 31-37, 1976.
- 64) Himori, N., Ono, H. and *Taira, N.*: Simultaneous assessment of effects of coronary vasodilators on the coronary blood flow and the myocardial contractility by using the blood-perfused canine papillary muscle. *Jpn. J. Pharmacol.* 26: 427-435, 1976.

- 65) Narimatsu, A. and *Taira, N.*: Effects on atrio-ventricular conduction of calcium-antagonistic coronary vasodilators, local anaesthetics and quinidine injected into the posterior and the anterior septal artery of the atrio-ventricular node preparation of the dog. *Naunyn-Schmiedeberg's Arch. Pharmacol.* 294: 169-177, 1976.
- 66) Ono, H., *Taira, N.* and Hashimoto, K.: Behavioural and vegetative effects of dibutyryl cyclic AMP on conscious dogs. *Neuropharmacology* 15: 571-575, 1976.
- 67) *Taira, N.*, Himori, N. and Imai, Y.: Inhibition of excitation-contraction coupling in the ventricular myocardium of the dog by SKF 24260. *Clin. Exp. Pharmacol. Physiol.* 3: 567-574, 1976.
- 68) Imai, Y., Himori, N. and *Taira, N.*: Cardiohemodynamic effects of SK&F 24260, D 600, diltiazem, dilazep, and trimetazidine in the dog. *Jpn. Heart J.* 18: 120-131, 1977.
- 69) Narimatsu, A. and *Taira, N.*: Positive dromotropic effect of dibutyryl cyclic adenosine 3',5'-monophosphate on the atrioventricular node. *Eur. J. Pharmacol.* 42: 71-78, 1977.
- 70) *Taira, N.*, Yabuuchi, Y. and Yamashita, S.: Profile of β -adrenoceptors in femoral, superior mesenteric and renal vascular beds of dogs. *Br. J. Pharmacol.* 59: 577-583, 1977.
- 71) Ono, H., Himori, N. and *Taira, N.*: Chronotropic effects of coronary vasodilators as assessed in the isolated, blood-perfused sino-atrial node preparation of the dog. *Tohoku J. Exp. Med.* 121: 383-390, 1977.
- 72) Himori, N., Imai, Y. and *Taira, N.*: Positive inotropic action of glyceryl trinitrate as observed in the blood-perfused papillary muscle preparation of the dog heart. *Clin. Exp. Pharmacol. Physiol.* 4: 257-262, 1977.
- 73) Himori, N. and *Taira, N.*: Assessment of the selectivity of OPC-2009, a new β_2 -adrenoceptor stimulant, by the use of the blood-perfused trachea *in situ* and of the isolated blood-perfused papillary muscle of the dog. *Br. J. Pharmacol.* 61: 9-17, 1977.
- 74) Narimatsu, A., Satoh, K. and *Taira, N.*: Assessment of the coronary vasodilator action of SK&F 24260 in the dog. *Clin. Exp. Pharmacol. Physiol.* 5: 107-115, 1978.
- 75) Endoh, M., Yanagisawa, T. and *Taira, N.*: Effects of calcium-antagonistic coronary vasodilators, nifedipine and verapamil, on ventricular automaticity of the dog. *Naunyn-Schmiedeberg's Arch. Pharmacol.* 302: 235-238, 1978.

- 76) Yanagisawa, T., Satoh, K. and *Taira, N.*: Excitation of autonomic nerves by 4-aminopyridine in the isolated, blood-perfused sino-atrial node preparation of the dog. *Eur. J. Pharmacol.* 49: 189-192, 1978.
- 77) *Taira, N.*, Iijima, T., Narimatsu, A., Satoh, K. and Yanagisawa, T.: Effects on atrio-ventricular conduction of propranolol, pindolol and carteolol in the dog heart *in situ* as assessed by automated devices. *Jpn. J. Pharmacol.* 28: 473-483, 1978.
- 78) Shimizu, T. and *Taira, N.*: Vasodilatation by prostaglandin $F_{2\alpha}$ in the canine tongue through a parasympathetic mechanism. *Br. J. Pharmacol.* 63: 567-574, 1978.
- 79) Imai, Y., Satoh, K. and *Taira, N.*: Role of the peripheral vasculature in changes in venous return caused by isoproterenol, norepinephrine, and methoxamine in anesthetized dogs. *Circ. Res.* 43: 553-561, 1978.
- 80) Hiwatari, M. and *Taira, N.*: Differential effects of D 600 on release of catecholamines by acetylcholine, histamine, tyramine and by cyclic AMP from canine adrenal medulla. *Jpn. J. Pharmacol.* 28: 671-680, 1978.
- 81) Himori, N. and *Taira, N.*: Pharmacological analysis of histamine receptors in musculature and vasculature of the dog trachea *in situ*. *Br. J. Pharmacol.* 64: 553-558, 1978.
- 82) Yanagisawa, T., Satoh, K. and *Taira, N.*: Simultaneous assessment of the effects of trimetazidine on the myocardium and coronary vasculature of the dog. *Arch. Int. Pharmacodyn. Ther.* 237: 316-329, 1979.
- 83) *Taira, N.*, Narimatsu, A., Satoh, K., Himori, N., Imai, Y. and Hiwatari, M.: Cardiovascular actions of niludipine (BAY a 7168), a new dihydropyridine vasodilator, in the dog. *Arzneim.-Forsch.* 29 (I): 246-255, 1979.
- 84) Hiwatari, M., Satoh, K. and *Taira, N.*: Antihypertensive effect of nifedipine on conscious renal hypertensive dogs. *Arzneim.-Forsch.* 29 (I): 256-260, 1979.
- 85) Shimizu, T. and *Taira, N.*: Assessment of the effects of vasoactive intestinal peptide (VIP) on blood flow through and salivation of the dog salivary gland in comparison with those of secretin, glucagon and acetylcholine. *Br. J. Pharmacol.* 65: 683-687, 1979.
- 86) *Taira, N.*, Satoh, K., Yanagisawa, T., Imai, Y. and Hiwatari, M.: Pharmacological profile of a new coronary vasodilator drug, 2-nicotin-amidoethyl nitrate (SG-75). *Clin. Exp. Pharmacol. Physiol.* 6: 301-315, 1979.

- 87) Yanagisawa, T. and *Taira, N.*: Positive inotropic effect of 4-aminopyridine on dog ventricular muscle. Naunyn-Schmiedeberg's Arch. Pharmacol. 307: 207-212, 1979.
- 88) Nakano, T. and *Taira, N.*: Potentiation of the algogenic action of bradykinin by an inhibitor of angiotensin I converting enzyme, captopril (SQ 14,225). Jpn. J. Pharmacol. 29: 563-571, 1979.
- 89) Satoh, K., Yanagisawa, T. and *Taira, N.*: Effects on atrioventricular conduction and blood flow of enantiomers of verapamil and of tetrodotoxin injected into the posterior and the anterior septal artery of the atrioventricular node preparation of the dog. Naunyn-Schmiedeberg's Arch. Pharmacol. 308: 89-98, 1979.
- 90) Hiwatari, M. and *Taira, N.*: Antihypertensive effect of niludipine (Bay a 7168) on conscious renal-hypertensive dogs. Arzneim.-Forsch. 29 (II): 1373-1376, 1979.
- 91) Yanagisawa, T., Satoh, K. and *Taira, N.*: Circumstantial evidence for increased potassium conductance of membrane of cardiac muscle by 2-nicotinamidoethyl nitrate (SG-75). Jpn. J. Pharmacol. 29: 687-694, 1979.
- 92) Himori, N., Ishimori, T. and *Taira, N.*: A further study on antihypertensive action of β -adrenoceptor blocking agents in conscious, renal hypertensive dogs. Arch. Int. Pharmacodyn. Ther. 242: 115-127, 1979.
- 93) Endoh, M., Yanagisawa, T. and *Taira, N.*: Effects of 2-nicotinamidethyl nitrate (SG-75), a new antianginal drug, on the cyclic AMP phosphodiesterase activity. Tohoku J. Exp. Med. 130: 199-201, 1980.
- 94) Himori, N. and *Taira, N.*: Differential effects of the calcium-antagonistic vasodilators, nifedipine and verapamil, on the tracheal musculature and vasculature of the dog. Br. J. Pharmacol. 68: 595-597, 1980.
- 95) Shimizu, T. and *Taira, N.*: Pharmacological analysis of salivary and blood flow responses to histamine of the submandibular gland of the dog. Br. J. Pharmacol. 68: 651-661, 1980.
- 96) Satoh, K., Yanagisawa, T. and *Taira, N.*: Mechanisms underlying the cardiovascular action of a new dihydropyridine vasodilator, YC-93. Clin. Exp. Pharmacol. Physiol. 7: 249-262, 1980.
- 97) Satoh, K., Yanagisawa, T. and *Taira, N.*: Coronary vasodilator and cardiac effects of optical isomers of verapamil in the dog. J. Cardiovasc. Pharmacol. 2: 309-318, 1980.

- 98) Yanagisawa, T. and *Taira, N.*: Effect of 2-nicotinamidethyl nitrate (SG-75) on the membrane potential of left atrial muscle fibres of the dog. Increase in potassium conductance. *Naunyn-Schmiedeberg's Arch. Pharmacol.* 312: 69-76, 1980.
- 99) Motomura, S., Iijima, T. and *Taira, N.*: Cholinergic intervention in intracardiac autonomic nerves in atrioventricular junctional area. *Am. J. Physiol.* 239 (Heart Circ. Physiol. 8): H181-H188, 1980.
- 100) Satoh, K., Yamashita, S., Endoh, M. and *Taira, N.*: Mechanism of cardiovascular action of trapidil. *Arzneim.-Forsch.* 30 (II): 1264-1268, 1980.
- 101) Yanagisawa, T., Iijima, T. and *Taira, N.*: Effects of trapidil on electrical activities of canine ventricular muscle. *Arzneim.-Forsch.* 30 (II): 1268-1271, 1980.
- 102) *Taira, N.*, Imai, Y. and Hiwatari, M.: Differential effects of nitroglycerin, trimetazidine, verapamil and SK&F 24260 on venous return as revealed by the open-loop method in the dog. *Jpn. J. Pharmacol.* 30: 449-461, 1980.
- 103) Endoh, M., Yanagisawa, T. and *Taira, N.*: Dissociation of cyclic AMP and contractile responses to isoprenaline: effects of a dihydropyridine derivative, nicardipine (YC-93), on canine ventricular muscle. *Eur. J. Pharmacol.* 67: 225-233, 1980.
- 104) Hashimoto, K., Motomura, S., Iijima, T. and *Taira, N.*: Effects of ouabain on atrioventricular conduction: Simultaneous observation in the isolated AV node preparation and the in situ heart of the dog. *Tohoku J. Exp. Med.* 132: 245-259, 1980.
- 105) Motomura, S. and *Taira, N.*: Differential effects of organic slow inward current inhibitors, verapamil and nifedipine, on rate of atrioventricular rhythm and supraventricular tachycardia in the canine isolated, blood-perfused AV node preparation. *Naunyn-Schmiedeberg's Arch. Pharmacol.* 315: 241-248, 1981.
- 106) Maruyama, M., Satoh, K. and *Taira, N.*: Effects on glyceryl trinitrate and SIN-1A on musculature and vasculature of the dog trachea *in situ*. *Br. J. Pharmacol.* 73: 9-11, 1981.
- 107) Shimizu, T., Nunoki, K. and *Taira, N.*: Characterization of neuronal and vascular histamine receptors mediating the salivary and vasodilator responses to histamine of the dog submandibular gland. *Tohoku J. Exp. Med.* 134: 1-8, 1981.

- 108) Yanagisawa, T. and *Taira, N.*: Effect of 2-nicotinamidethyl nitrate (SG-75) on membrane potentials of canine Purkinje fibers. *Jpn. J. Pharmacol.* 31: 409-417, 1981.
- 109) Yamashita, S., Motomura, S. and *Taira, N.*: Cardiac effects of amiloride in the dog. *J. Cardiovasc. Pharmacol.* 3: 704-715, 1981.
- 110) Satoh, K., Narimatsu, A. and *Taira, N.*: Effects of antiarrhythmic drugs on AV nodal and intraventricular conduction as assessed in the isolated, blood-perfused AV node preparation of the dog. *J. Cardiovasc. Pharmacol.* 3: 753-768, 1981.
- 111) Maruyama, M., Satoh, K. and *Taira, N.*: Effects of trapidil on musculature and vasculature of the dog trachea *in situ*. *Arch. Int. Pharmacodyn. Ther.* 252: 307-312, 1981.
- 112) Motomura, S., Satoh, K., Yamashita, S. and *Taira, N.*: Characterization of cardiovascular actions of CS-611, a new coronary dilator, in the dog. *Arzneim.-Forsch.* 31 (II): 1238-1244, 1981.
- 113) Tokudome, T. and *Taira, N.*: Characterization of β -adrenoceptors in the dog saphenous vein. *Jpn. J. Pharmacol.* 31: 731-736, 1981.
- 114) Maruyama, M., Satoh, K. and *Taira, N.*: Effects of amrinone on the tracheal musculature and vasculature of the dog. *Jpn. J. Pharmacol.* 31: 1095-1097, 1981.
- 115) Endoh, M., Satoh, K. and *Taira, N.*: α -Adrenoceptors mediating the positive inotropic effect of phenylephrine in the right ventricular muscle of the monkey (*Macaca fuscata*). *Naunyn-Schmiedeberg's Arch. Pharmacol.* 318: 370-373, 1982.
- 116) Shimizu, T., Nunoki, K. and *Taira, N.*: Further evidence for postganglionic actions of prostaglandin $F_{2\alpha}$ and histamine in the submandibular gland of the dog. *Tohoku J. Exp. Med.* 137: 59-63, 1982.
- 117) Endoh, M., Yamashita, S. and *Taira, N.*: Positive inotropic effect of amrinone in relation to cyclic nucleotide metabolism in the canine ventricular muscle. *J. Pharmacol. Exp. Ther.* 221: 775-783, 1982.
- 118) Maruyama, M., Satoh, K. and *Taira, N.*: Effects of nicorandil and its congeners on musculature and vasculature of the dog trachea *in situ*. *Arch. Int. Pharmacodyn. Ther.* 258: 260-266, 1982.

- 119) Satoh, K., Yamashita, S., Maruyama, M. and *Taira, N.*: Comparison of the responses of the simian and canine coronary circulations to autonomic drugs. *J. Cardiovasc. Pharmacol.* 4: 820-828, 1982.
- 120) Satoh, K., Maruyama, M. and *Taira, N.*: The improvement of cardiac performance by amrinone, a new cardiotonic drug, in an experimental failing heart preparation of the dog. *Jpn. Heart J.* 23: 975-980, 1982.
- 121) Endoh, M., Maruyama, M. and *Taira, N.*: Modification by islet-activating protein of direct and indirect inhibitory actions of adenosine on rat atrial contraction in relation to cyclic nucleotide metabolism. *J. Cardiovasc. Pharmacol.* 5: 131-142, 1983.
- 122) *Taira, N.*, Satoh, K., Maruyama, M. and Yamashita, S.: Sustained coronary constriction and its antagonism by calcium-blocking agents in monkeys and baboons. *Circ. Res.* 52 (Suppl. I): 40-46, 1983.
- 123) Satoh, K., Kawada, M. and *Taira, N.*: Improvement by trapidil of cardiac function of the dog heart-lung preparation depressed by pentobarbital. *Jpn. Heart J.* 24: 399-405, 1983.
- 124) *Taira, N.*, Kawada, M. and Satoh, K.: Cardiac versus coronary vasodilator actions of KB-944, a new calcium antagonist, assessed in isolated, blood-perfused heart preparations of dogs. *J. Cardiovasc. Pharmacol.* 5: 349-356, 1983.
- 125) Endoh, M. and *Taira, N.*: Relationship between relaxation and cyclic GMP formation caused by nicorandil in canine mesenteric artery. *Naunyn-Schmiedeberg's Arch. Pharmacol.* 322: 319-321, 1983.
- 126) Kawada, M., Satoh, K. and *Taira, N.*: Profile of coronary vasodilator and cardiac actions of bepridil revealed by use of isolated, blood-perfused heart preparations of the dogs. *J. Cardiovasc. Pharmacol.* 5: 604-612, 1983.
- 127) Iijima, T. and *Taira, N.*: Effects of manganese ions and diltiazem on the spontaneous action potential of the canine atrio-ventricular node cell. *J. Mol. Cell. Cardiol.* 15: 863-866, 1983.
- 128) Kawada, M., Satoh, K. and *Taira, N.*: Analyses of the cardiac action of the bradycardic agent, AQ-A 39, by use of isolated, blood-perfused dog-heart preparations. *J. Pharmacol. Exp. Ther.* 228: 484-490, 1984.
- 129) *Taira, N.*, Endoh, M., Iijima, T., Satoh, K., Yanagisawa, T., Yamashita, S., Maruyama, M., Kawada, M., Morita, T. and Wada, Y.: Mode and mechanism of action of 3,4-dihydro-6-[4-(3,4-dimethoxybenzoyl)-1-piperazinyl]-2(1H)-quinolinone (OPC-8212), a novel positive inotropic drug, on the dog heart. *Arzneim.-Forsch.* 34 (I): 347-355, 1984.

- 130) Satoh, K., Maruyama, M., Yamashita, S. and *Taira, N.*: Nicorandil releases acetylcholine-induced sustained coronary arterial constriction in monkeys and baboons. *Jpn. Heart J.* 25: 267-273, 1984.
- 131) Satoh, K., Kawada, M., Wada, Y. and *Taira, N.*: Cardiovascular actions of the dihydropyridine calcium antagonist nimodipine in the dog. *Arzneim.-Forsch.* 34 (I): 563-568, 1984.
- 132) Morita, T., Satoh, K. and *Taira, N.*: Improvement of cardiac performance by AR-L 115 BS, a new cardiotonic drug, in the experimental failing dog heart. *Jpn. Heart J.* 25: 439-445, 1984.
- 133) Morita, T., Satoh, K. and *Taira, N.*: Comparison of the cardiovascular and biochemical profiles of a new positive inotropic drug, AR-L 115 BS, with those of theophylline. *Jpn. Heart J.* 25: 447-460, 1984.
- 134) Satoh, K., Wada, Y. and *Taira, N.*: Differential effects of Bay k 8644, a presumed calcium channel activator, on sinoatrial nodal and ventricular automaticity of the dog heart. *Naunyn-Schmiedeberg's Arch. Pharmacol.* 326: 190-192, 1984.
- 135) Wada, Y., Satoh, K. and *Taira, N.*: Coronary vasodilator compared to cardiac effects of PY 108-068, a new dihydropyridine vasodilator, on isolated, blood-perfused dog-heart preparations. *J. Cardiovasc. Pharmacol.* 6: 881-887, 1984.
- 136) Kawada, M., Satoh, K. and *Taira, N.*: Selectivity of alinidine, a bradycardic agent, for SA nodal automaticity versus other cardiac activities in isolated, blood-perfused dog-heart preparations. *Arch. Int. Pharmacodyn. Ther.* 272: 88-102, 1984.
- 137) Yanagisawa, T., Endoh, M. and *Taira, N.*: Involvement of cyclic AMP in the positive inotropic effect of OPC-8212, a new cardiotonic agent, on the canine ventricular muscle. *Jpn. J. Pharmacol.* 36: 379-388, 1984.
- 138) Iijima, T., Yanagisawa, T. and *Taira, N.*: Increase in the slow inward current by intracellularly applied nifedipine and nicardipine in single ventricular cells of the guinea-pig heart. *J. Mol. Cell. Cardiol.* 16: 1173-1177, 1984.
- 139) *Taira, N.*, Wada, Y. and Satoh, K.: Is antagonism by Bay k 8644 of the negative dromotropic effect of nifedipine pharmacological? *J. Pharmacol. Exp. Ther.* 232: 244-250, 1985.
- 140) Ishii, K., Yanagisawa, T., Satoh, K. and *Taira, N.*: Abolition of spontaneous rhythmic contractions of isolated monkey coronary arteries by diltiazem, nifedipine, verapamil and nicorandil but not by nitroglycerin. *Tohoku J. Exp. Med.* 145: 85-90, 1985.

- 141) Wada, Y., Satoh, K. and *Taira, N.*: Separation of the coronary vasodilator from the cardiac effects of PN 200-110, a new dihydropyridine calcium antagonist, in the dog heart. *J. Cardiovasc. Pharmacol.* 7: 190-196, 1985.
- 142) Ishii, K., *Taira, N.* and Yanagisawa, T.: Differential antagonism by Bay k 8644, a dihydropyridine calcium agonist, of the negative inotropic effects of nifedipine, verapamil, diltiazem and manganese ions in canine ventricular muscle. *Br. J. Pharmacol.* 84: 577-584, 1985.
- 143) Wada, Y., Satoh, K. and *Taira, N.*: Cardiovascular profile of Bay k 8644, a presumed calcium channel activator, in the dog. *Naunyn-Schmiedeberg's Arch. Pharmacol.* 328: 382-387, 1985.
- 144) Satoh, K., Kawada, M., Wada, Y. and *Taira, N.*: Nicardipine releases sustained coronary arterial constriction induced by acetylcholine in the rhesus monkey. *Heart Vessels.* 1: 74-77, 1985.
- 145) Satoh, K., Nunoki, K., Goto, T. and *Taira, N.*: The mechanism underlying the vasodilator action of bunitrolol: Contribution of α_1 -adrenoceptor blocking action. *J. Cardiovasc. Pharmacol.* 7: 508-515, 1985.
- 146) Endoh, M., Yanagisawa, T., Morita, T. and *Taira, N.*: Differential effects of sulmazole (AR-L 115 BS) on contractile force and cyclic AMP levels in canine ventricular muscle: Comparison with MDL 17, 043. *J. Pharmacol. Exp. Ther.* 234: 267-273, 1985.
- 147) Goto, T., Satoh, K. and *Taira, N.*: Bay K 8644, a dihydropyridine calcium agonist, augments vasoconstrictor responses to endogenous and exogenous noradrenaline in the peripheral vasculature of the dog. *Br. J. Pharmacol.* 85: 913-916, 1985.
- 148) Iijima, T. and *Taira, N.*: Abolition by nicorandil of spontaneous action potentials of the small atrio-ventricular node preparation of the dog. *Tohoku J. Exp. Med.* 146: 401-403, 1985.
- 149) Iijima, T., Morita, T. and *Taira, N.*: Effects of a new calcium entry blocking agent, KB-944, on the membrane potential and currents of mammalian ventricular muscle cells. *Arch. Int. Pharmacodyn. Ther.* 277: 77-82, 1985.
- 150) Morita, T., Maniwa, T., Satoh, K. and *Taira, N.*: Undifferentiated effects of calcium antagonists on pressor responses to selective α -1 and α -2 adrenoceptor agonists in anesthetized, spinal dogs. *J. Pharmacol. Exp. Ther.* 234: 728-734, 1985.

- 151) Nunoki, K., Goto, T., Satoh, K. and *Taira, N.*: Improvement by denopamine (TA-064) of pentobarbital-induced cardiac failure in the dog heart-lung preparation. *Heart Vessels* 1: 216-219, 1985.
- 152) Goto, T., Nunoki, K., Satoh, K. and *Taira, N.*: Amelioration of heart failure by MDL 17043 and MDL 19205, novel positive inotropic drugs, in dog heart-lung preparations. *Jpn. Heart J.* 26: 1011-1018, 1985.
- 153) Satoh, K., Wada, Y. and *Taira, N.*: Comparative study of coronary vasodilator and cardiac effects of nitrendipine by use of isolated, blood-perfused dog heart preparations. *Arzneim.-Forsch.* 36 (I): 35-39, 1986.
- 154) Wada, Y. and *Taira, N.*: Comparative study of MDL 17043 and MDL 19205, new positive inotropic agents, by use of isolated, blood-perfused dog-heart preparations. *Heart Vessels* 2: 29-35, 1986.
- 155) Takahashi, K. and *Taira, N.*: Cardiac versus coronary vasodilator actions of isobutylmethylxanthine in dogs: Comparison with theophylline. *Jpn. J. Pharmacol.* 41: 23-32, 1986.
- 156) Ishii, K., Sato, Y. and *Taira, N.*: Similarity and dissimilarity of the vasoconstrictor effects of Bay K 8644 on coronary, femoral, mesenteric and renal circulations of dogs. *Br. J. Pharmacol.* 88: 369-377, 1986.
- 157) Takahashi, K. and *Taira, N.*: Cardiovascular profiles of the calcium antagonists nisoldipine and nifedipine in blood-perfused dog heart. *Arzneim.-Forsch.* 36 (II): 1054-1059, 1986.
- 158) Kawada, M., Satoh, K. and *Taira, N.*: Cardiohemodynamic effects of nipradilol (K-351) in the dog: Comparison with propranolol, nadolol and prazosin. *Jpn. J. Pharmacol.* 42: 9-18, 1986.
- 159) Sato, Y., Wada, Y. and *Taira, N.*: Comparative study of cardiovascular profiles of milrinone and amrinone by use of isolated, blood-perfused dog heart preparations. *Heart Vessels* 2: 213-220, 1986.
- 160) Iijima, T. and *Taira, N.*: Modification by nicorandil and carbachol of the inward calcium current in single cells of the guinea-pig heart. *Tohoku J. Exp. Med.* 150: 475-479, 1986.
- 161) Imai, Y., Abe, K., Sasaki, S., Minami, N., Seino, M., Yoshinaga, K. and *Taira, N.*: The mechanism of centrally mediated cardiovascular actions of the three structurally different calcium antagonists, verapamil, diltiazem and nifedipine, in rats. *Tohoku J. Exp. Med.* 151: 65-80, 1987.

- 162) Iijima, T. and *Taira, N.*: Membrane current changes responsible for the positive inotropic effect of OPC-8212, a new positive inotropic agent, in single ventricular cells of the guinea pig heart. *J. Pharmacol. Exp. Ther.* 240: 657-662, 1987.
- 163) Takahashi, K., Endoh, M. and *Taira, N.*: Inotropic versus chronotropic, dromotropic, and coronary vasodilator actions of DPI 201-106, a novel positive inotropic agent, in the dog heart. *J. Cardiovasc. Pharmacol.* 9: 451-460, 1987.
- 164) Iijima, T., Endoh, M. and *Taira, N.*: Differential effects of isoproterenol on the canine atrial action potential in the presence of carbachol or nicorandil. *Jpn. J. Pharmacol.* 44: 218-221, 1987.
- 165) Hosono, M. and *Taira, N.*: Coronary vasodilator versus cardiac effects of MCI-176, a novel quinazolinone calcium antagonist, in the dog heart. *J. Cardiovasc. Pharmacol.* 9: 633-640, 1987.
- 166) Yanagisawa, T., Ishii, K. and *Taira, N.*: Antitachycardiac effect of OPC-8212, a novel cardiotonic agent, on tachycardiac responses of guinea pig isolated right atria to isoproterenol and histamine. *J. Cardiovasc. Pharmacol.* 10: 47-54, 1987.
- 167) Hashimoto, H., Ishii, K., Satoh, K. and *Taira, N.*: Comparison of vasodilator effects of DN-9693, a selective inhibitor of cyclic AMP phosphodiesterase, and isobutylmethylxanthine, a non-selective one, in dogs. *Jpn. J. Pharmacol.* 44: 405-412, 1987.
- 168) Katayama, S., Narimatsu, A., Suzuki, R., Iijima, T. and *Taira, N.*: Changes in membrane potentials and currents of ventricular cells of the guinea pig heart by a new cardiotonic drug, MCI-154. *Jpn. J. Pharmacol.* 44: 481-488, 1987.
- 169) Iijima, T. and *Taira, N.*: Pinacidil increases the background potassium current in single ventricular cells. *Eur. J. Pharmacol.* 141: 139-141, 1987.
- 170) *Taira, N.*, Takahashi, K. and Hosono, M.: Effect of DHP-218, a novel dihydropyridine phosphonate, on atrioventricular nodal conductivity compared with its vascular effect in dogs. *J. Cardiovasc. Pharmacol.* 10: 274-279, 1987.
- 171) Kitada, Y., Narimatsu, A., Suzuki, R., Endoh, M. and *Taira, N.*: Does the positive inotropic action of a novel cardiotonic agent, MCI-154, involve mechanisms other than cyclic AMP? *J. Pharmacol. Exp. Ther.* 243: 639-645, 1987.

- 172) Hosono, M. and *Taira, N.*: Cardiac and coronary vasodilator effects of the novel cardiotonic agent, MCI-154, assessed in isolated, blood-perfused dog heart preparations. *J. Cardiovasc. Pharmacol.* 10: 692-698, 1987.
- 173) Imagawa, J., Satoh, K. and *Taira, N.*: Cardiac and coronary vasodilator profile of pimobendan, a new cardiotonic drug, revealed by use of isolated, blood-perfused dog heart preparations. *Heart Vessels* 3: 182-189, 1987.
- 174) Sato, Y., Ishii, K. and *Taira, N.*: Differential antagonism by Bay K 8644 of vasodilator effects of nifedipine, diltiazem, nicorandil and nitroglycerin in dog femoral circulation. *Br. J. Pharmacol.* 93: 146-150, 1988.
- 175) Iijima, T., Takahashi, K. and *Taira, N.*: Voltage- and use-dependent block of the inward calcium current by MCI-176, a new non-dihydropyridine calcium antagonist, in canine ventricular muscles and single ventricular cells of the guinea-pig. *Jpn. J. Pharmacol.* 46: 155-164, 1988.
- 176) Satoh, K., Nunoki, K., Goto, T., Hosono, H., Hashimoto H., Sato, Y. and *Taira, N.*: Improvement of pentobarbital-induced heart failure by MCI-154, a novel and potent cardiotonic agent, in the dog heart-lung preparation. *Jpn. J. Pharmacol.* 47: 189-195, 1988.
- 177) Hosono, M., Satoh, K. and *Taira, N.*: T-1583 and forskolin are similar in their cardiac effects and dissimilar in their vascular effects. *Cardiovasc. Drugs Ther.* 2: 245-253, 1988.
- 178) Gotanda, K., Yanagisawa, T., Satoh, K. and *Taira, N.*: Are the cardiovascular effects of gentamicin similar to those of calcium antagonists? *Jpn. J. Pharmacol.* 47: 217-227, 1988.
- 179) Gotanda, K., Satoh, K. and *Taira, N.*: Is the cardiovascular profile of BRL 34915 characteristic of potassium channel activators? *J. Cardiovasc. Pharmacol.* 12: 239-246, 1988.
- 180) Umekawa, H., Yamakawa, K., Nunoki, K., *Taira, N.*, Tanaka, K. and Hidaka, H.: Inhibition of calmodulin function by CV-159, a novel dihydropyridine compound. *Biochem. Pharmacol.* 37: 3377-3381, 1988.
- 181) Yokoyama, H., Yanagisawa, T. and *Taira, N.*: Details of mode and mechanism of action of denopamine, a new orally active cardiotonic agent with affinity for β_1 -receptors. *J. Cardiovasc. Pharmacol.* 12: 323-331, 1988.
- 182) Gotanda, K., Satoh, K. and *Taira, N.*: Cardiovascular profile of AN-132, a new diamine derivative antiarrhythmic agent, revealed in the isolated, blood-perfused dog heart. *Tohoku J. Exp. Med.* 156: 65-78, 1988.

- 183) Yanagisawa, T., Hashimoto, H. and *Taira, N.*: The negative inotropic effect of nicorandil is independent of cyclic GMP changes: A comparison with pinacidil and cromakalim in canine atrial muscle. *Br. J. Pharmacol.* 95: 393-398, 1988.
- 184) Yokoyama, H., Imagawa, J., Satoh, K., *Taira, N.* and Tamahashi, N.: Isolated dog hearts prepared in cold Tyrode solution and reperfused with arterial blood are functionally and ultrastructurally normal. *Tohoku J. Exp. Med.* 156: 121-134, 1988.
- 185) Sato, Y., Iijima, T. and *Taira, N.*: Electrophysiological effects of the new antiarrhythmic drug 3-(diisopropylaminoethylamino)-2',6'-dimethylpropionanilide on dog cardiac muscle. *Arzneim.-Forsch.* 39 (I): 56-59, 1989.
- 186) Yanagisawa, T., Ishii, K., Hashimoto, H. and *Taira, N.*: Differential coupling to positive inotropic responses of cyclic AMP produced by stimulation of β_1 - and β_2 -adrenergic receptors. *J. Cardiovasc. Pharmacol.* 13: 64-75, 1989.
- 187) Imagawa, J., Satoh, K. and *Taira, N.*: Coronary vasodilator and cardiac effects of NKY-722, a novel hydrophilic 1,4-dihydropyridine derivative, in the blood-perfused dog heart. *Cardiovasc. Drugs Ther.* 3: 81-90, 1989.
- 188) Hashimoto, H., Yanagisawa, T. and *Taira, N.*: Differential antagonism of the negative inotropic effect of gentamicin by calcium ions, Bay K 8644 and isoprenaline in canine ventricular muscle: Comparison with cobalt ions. *Br. J. Pharmacol.* 96: 906-912, 1989.
- 189) Iijima, T. and *Taira, N.*: β_2 -Adrenoceptor-mediated increase in the slow inward calcium current in atrial cells. *Eur. J. Pharmacol.* 163: 357-360, 1989.
- 190) Yanagisawa, T., Hashimoto, H. and *Taira, N.*: Interaction of potassium channel openers and blockers in canine atrial muscle. *Br. J. Pharmacol.* 97: 753-762, 1989.
- 191) Gotanda, K., Yokoyama, H., Satoh, K. and *Taira, N.*: Cardiohemodynamic effects of cromakalim and pinacidil, potassium-channel openers, in the dog, special reference to venous return. *Cardiovasc. Drugs Ther.* 3: 507-515, 1989.
- 192) Koga, Y., Sodeyama, N., Satoh, S., Teshigawara, T., Yanagisawa, T., *Taira, N.* and Endoh, M.: Relaxation of airway smooth muscle induced by potassium in the presence of Ca-antagonists. *Jpn. J. Pharmacol.* 50: 387-396, 1989.

- 193) Yanagisawa, T., Kawada, M. and *Taira, N.*: Nitroglycerin relaxes canine coronary arterial smooth muscle without reducing intracellular Ca^{2+} concentrations measured with fura-2. *Br. J. Pharmacol.* 98: 469–482, 1989.
- 194) Yokoyama, H., Ishii, K., Yanagisawa, T. and *Taira, N.*: An explanation of bell-shaped, dose-positive inotropic effect curves for denopamine in canine right ventricular muscle. *J. Cardiovasc. Pharmacol.* 14: 570–576, 1989.
- 195) Iijima, T. and *Taira, N.*: Mechanisms of β -adrenoceptor mediated increase in delayed rectifier potassium current. *Tohoku J. Exp. Med.* 159: 277–281, 1989.
- 196) Mori, T., Yanagisawa, T. and *Taira, N.*: Histamine increases vascular tone and intracellular calcium level using both intracellular and extracellular calcium in porcine coronary arteries. *Jpn. J. Pharmacol.* 52: 263–271, 1990.
- 197) Mori, T., Yanagisawa, T. and *Taira, N.*: Phorbol 12,13-dibutyrate increases vascular tone but has a dual action on intracellular calcium levels in porcine coronary arteries. *Naunyn-Schmiedeberg's Arch. Pharmacol.* 341: 251–255, 1990.
- 198) Yamada, H., Yoneyama, F., Satoh, K. and *Taira, N.*: Specific but differential antagonism by glibenclamide of the vasodepressor effects of cromakalim and nicorandil in spinally-anaesthetized dogs. *Br. J. Pharmacol.* 100: 413–416, 1990.
- 199) Iijima, T., Imagawa, J. and *Taira, N.*: Differential modulation by *beta* adrenoceptors of inward calcium and delayed rectifier potassium current in single ventricular cells of guinea pig heart. *J. Pharmacol. Exp. Ther.* 254: 142–146, 1990.
- 200) Yoneyama, F., Satoh, K. and *Taira, N.*: Nicorandil increases coronary blood flow predominantly by K-channel opening mechanism. *Cardiovasc. Drugs Ther.* 4: 1119–1126, 1990.
- 201) Yanagisawa, T., Teshigawara, T., and *Taira, N.*: Cytoplasmic calcium and the relaxation of canine coronary arterial smooth muscle produced by cromakalim, pinacidil and nicorandil. *Br. J. Pharmacol.* 101: 157–165, 1990.
- 202) Satoh, E., Yanagisawa, T. and *Taira, N.*: Specific antagonism by glibenclamide of negative inotropic effects of potassium channel openers in canine atrial muscle. *Jpn. J. Pharmacol.* 54: 133–141, 1990.
- 203) Kurosawa, H., Satoh, E., Yanagisawa, T. and *Taira, N.*: Selective β_1 -receptor full agonists, T-0509 and T-1583, increase the force monophasically and cyclic AMP biphasically in canine ventricular muscle. *J. Cardiovasc. Pharmacol.* 16: 646–653, 1990.

- 204) Yoneyama, F., Yamada, H., Satoh, K. and *Taira, N.*: Cardiac versus coronary dilator effects of SD-3211, a new nondihydropyridine calcium antagonist, in isolated, blood-perfused dog hearts. *Cardiovasc. Drugs Ther.* 4: 1469–1476, 1990.
- 205) Satoh, K., Yamada H. and *Taira, N.*: Differential antagonism by glibenclamide of the relaxant effects of cromakalim, pinacidil and nicorandil on canine large coronary arteries. *Naunyn-Schmiedeberg's Arch. Pharmacol.* 343: 76–82, 1991.
- 206) Yamada, H., Yoneyama, F., Satoh, K. and *Taira, N.*: Comparison of the effects of the novel vasodilator FK409 with those of nitroglycerin in isolated coronary artery of the dog. *Br. J. Pharmacol.* 103: 1713–1718, 1991.
- 207) Okada, Y., Yanagisawa, T. and *Taira, N.*: An analysis of the nitrate-like and K-channel opening actions of KRN2391 in canine coronary arterial smooth muscle. *Br. J. Pharmacol.* 104: 829–838, 1991.
- 208) Yamada, H., Yoneyama, F., Satoh, K. and *Taira, N.*: Cardiohemodynamic effect of FK409, a novel highly potent nitrovasodilator, in anesthetized dogs. *Eur. J. Pharmacol.* 205: 81–83, 1991.
- 209) Satoh, K., Yamada, H., Yoneyama, F. and *Taira, N.*: The group at C2 of N-ethylnicotinamide determines the vasodilator potencies and mechanisms of action of nicorandil and its congeners in coronary arteries. *Naunyn-Schmiedeberg's Arch. Pharmacol.* 344: 589–595, 1991.
- 210) Kageyama, M., Mori, T., Yanagisawa, T. and *Taira, N.*: Is staurosporine a specific inhibitor of protein kinase C in intact porcine coronary arteries? *J. Pharmacol. Exp. Ther.* 259: 1019–1026, 1991.
- 211) Imagawa, J., Akima, M., Nabata, H. and *Taira, N.*: Spasmolytic action of nicorandil in canine conductive coronary arteries in vivo is not modified by glibenclamide. *J. Cardiovasc. Pharmacol.* 19: 108–114, 1992.
- 212) Yoneyama, F., Yamada, H., Satoh, K. and *Taira, N.*: Vasodepressor mechanisms of 2-(1-octynyl)-adenosine (YT-146), a selective adenosine A₂ receptor agonist, involve the opening of glibenclamide-sensitive K⁺ channels. *Eur. J. Pharmacol.* 213: 199–204, 1992.
- 213) Ishii, K., Nunoki, K., Murakoshi, H. and *Taira, N.*: Cloning and modulation by endothelin-1 of rat cardiac K channel. *Biochem. Biophys. Res. Commun.* 184: 1484–1489, 1992.

- 214) Kageyama, M., Yamada, H., Satoh, K. and *Taira, N.*: Effects of semotiadil (SD-3211), a benzothiazine calcium antagonist, on blood pressure and atrioventricular conductivity in anesthetized dogs. *Jpn. J. Pharmacol.* 59: 177–182, 1992.
- 215) Yamagishi, T., Yanagisawa, T. and *Taira, N.*: Ca^{2+} influx induced by the agonist U46619 is inhibited by hyperpolarization induced by the K^{+} channel opener cromakalim in canine coronary artery. *Jpn. J. Pharmacol.* 59: 291–299, 1992.
- 216) Okada, Y., Yanagisawa, T. and *Taira, N.*: E4080 has a dual action, as a K^{+} channel opener and a Ca^{2+} channel blocker, in canine coronary artery smooth muscle. *Eur. J. Pharmacol.* 218: 259–264, 1992.
- 217) Yamagishi, T., Yanagisawa, T. and *Taira, N.*: Activation of phospholipase C by the agonist U46619 is inhibited by cromakalim-induced hyperpolarization in porcine coronary artery. *Biochem. Biophys. Res. Commun.* 187: 1517–1522, 1992.
- 218) Okada, H., Ishii, K., Nunoki, K., Abe, T. and *Taira, N.*: Modulation of transient type K channel cloned from rat heart. *Biochem. Biophys. Res. Commun.* 189: 430–436, 1992.
- 219) Orito, K., Satoh, K. and *Taira, N.*: ATP-sensitive K^{+} channels are gradually recruited in the vasodepressor response to adenosine in spinally-anesthetized dogs. *Jpn. J. Pharmacol.* 60: 295–297, 1992.
- 220) Okada, Y., Yanagisawa, T. and *Taira, N.*: KCl-depolarization potentiates the Ca^{2+} sensitization by endothelin-1 in canine coronary artery. *Jpn. J. Pharmacol.* 60: 403–405, 1992.
- 221) Yamagishi, T., Yanagisawa, T. and *Taira, N.*: K^{+} channel openers, cromakalim and Ki4032, inhibit agonist-induced Ca^{2+} release in canine coronary artery. *Naunyn-Schmiedeberg's Arch. Pharmacol.* 346: 691–700, 1992.
- 222) Orito, K., Satoh, K. and *Taira, N.*: Involvement of ATP-sensitive K^{+} channels in the sustained coronary vasodilator response to adenosine in dogs. *Eur. J. Pharmacol.* 231: 183–189, 1993.
- 223) Tofukuji, M., Yanagisawa, T., Kurosawa, H., Ishii, K. and *Taira, N.*: ICI 147,798 causes insurmountable antagonism against T-0509, a selective β_1 -adrenoceptor agonist, but surmountable antagonism against isoproterenol. *J. Cardiovasc. Pharmacol.* 21: 323–331, 1993.

- 224) Satoh, K., Sato, Y., Imagawa, J. and *Taira, N.*: Improvement of cardiac performance by pimobendan, a new cardiotonic drug, in the experimental failing dog heart. *Jpn. Heart J.* 34: 213-219, 1993.
- 225) Yamagishi, T., Ishii, K. and *Taira, N.*: Absence of effects of class III antiarrhythmic agents on cloned cardiac K channels. *Jpn. J. Pharmacol.* 61: 371-373, 1993.
- 226) Okada, Y., Yanagisawa, T. and *Taira, N.*: BRL 38227 (levcromakalim)-induced hyperpolarization reduces the sensitivity to Ca^{2+} of contractile elements in canine coronary artery. *Naunyn-Schmiedeberg's Arch. Pharmacol.* 347: 438-444, 1993.
- 227) Furukawa, S., Satoh, K. and *Taira, N.*: Opening of ATP-sensitive K^+ channels responsible for adenosine A_2 receptor-mediated vasodepression does not involve a pertussis toxin-sensitive G protein. *Eur. J. Pharmacol.* 236: 255-262, 1993.
- 228) Kageyama, M., Yanagisawa, T. and *Taira, N.*: Calcitonin gene-related peptide relaxes porcine coronary arteries *via* cyclic AMP-dependent mechanisms, but not activation of ATP-sensitive potassium channels. *J. Pharmacol. Exp. Ther.* 265: 490-497, 1993.
- 229) Satoh, K., Mori, T., Yamada H. and *Taira, N.*: Nicorandil as a nitrate, and cromakalim as a potassium channel opener, dilate isolated porcine large coronary arteries in an agonist-nonselective manner. *Cardiovasc. Drugs Ther.* 7: 691-699, 1993.
- 230) Orito, K., Satoh, K. and *Taira, N.*: Cardiovascular profile of Ro 40-5967, a new nondihydropyridine calcium antagonist, delineated in isolated, blood-perfused dog heart. *J. Cardiovasc. Pharmacol.* 2: 293-299, 1993.
- 231) Okada, Y., Yanagisawa, T., Yamagishi, T. and *Taira, N.*: K^+ channel-opening action and KRN2391-induced reduction of Ca^{2+} sensitivity of arterial smooth muscle. *Arch. Int. Pharmacodyn. Ther.* 326: 33-51, 1993.
- 232) Ishii, K., Yamagishi, T. and *Taira, N.*: Cloning and functional expression of a cardiac inward rectifier K^+ channel. *FEBS Lett.* 338: 107-111, 1994.
- 233) Yamagishi, T., Yanagisawa, T., Satoh, K. and *Taira, N.*: Relaxant mechanisms of cyclic AMP-increasing agents in porcine coronary artery. *Eur. J. Pharmacol.* 251: 253-262, 1994.
- 234) Orito, K., Satoh, K. and *Taira, N.*: Vasodilator mechanisms of action of amrinone do not involve the opening of either ATP-sensitive or Ca^{2+} -activated K^+ channels. *Tohoku. J. Exp. Med.* 172: 163-173, 1994.

- 235) Satoh, K., Orito, K., Yoneyama, F. and *Taira, N.*: A further study of the vasodilator and negative inotropic mechanisms of action of nicorandil and its congeners in the canine heart. *Cardiovasc. Drugs Ther.* 8: 227-234, 1994.
- 236) Orito, K., Yanagisawa, T. and *Taira, N.*: A possibility that the ATP-sensitive potassium channel in coronary artery has a high-affinity internal binding site for tetraalkylammonium. *Jpn. J. Pharmacol.* 64: 297-301, 1994.
- 237) Murakoshi, H., Ishii, K., Nunoki, K. and *Taira, N.*: Receptor-mediated modulation of rat $K_{V1.2}$ in *Xenopus* oocytes. *Eur. J. Pharmacol. Mol. Pharmacol. Sec.* 268: 451-454, 1994.
- 238) Nunoki, K., Ishii, K., Okada, H., Yamagishi, T., Murakoshi, H. and *Taira, N.*: Hybrid potassium channels by tandem linkage of inactivating and non-inactivating subunits. *J. Biol. Chem.* 269: 24138-24142, 1994.

Review Articles

- 1) *Taira, N.*: The autonomic pharmacology of the bladder. *Annu. Rev. Pharmacol.* 12: 197-208, 1972.
- 2) Hashimoto, K., *Taira, N.*, Ono, H., Chiba, S., Hashimoto, K., Endoh, M., Kokubun, M., Kokubun, H., Iijima, T., Kimura, T., Kubota, K. and Oguro, K.: Nifedipine, basis of its pharmacological effect. In: 1st International Nifedipine >> Adalat® << Symposium — New Therapy of Ischemic Heart Disease, ed. by K. Hashimoto, E. Kimura and T. Kobayashi, pp. 11-22, University of Tokyo Press, Tokyo, 1975.
- 3) *Taira, N.*: Effects of diltiazem and other calcium-antagonists on cardiac functions and coronary blood flow as assessed in blood-perfused dog-heart preparations. In: New Drug Therapy with a Calcium Antagonist — Diltiazem Hakone Symposium '78, ed. by R. J. Bing, pp. 91-103, Excerpta Medica, Amsterdam — Princeton, 1979.
- 4) Yanagisawa, T. and *Taira, N.*: Effects of 4-aminopyridine on cardiac muscle. In: *Adv. Biosci.* Vol. 35: Aminopyridines and Similarly Acting Drugs: Effects on Nerves, Muscles and Synapses, ed. by P. Lechat, S. Thesleff and W. C. Bowman, pp. 261-274, Pergamon Press, Oxford — New York, 1982.
- 5) *Taira, N.*, Satoh, K., Maruyama, M. and Yamashita, S.: Acetylcholine-induced sustained coronary constriction and its antagonism by calcium antagonists in monkeys and baboons. In: 5th International Adalat® Symposium — New Therapy of Ischaemic Heart Disease and Hypertension, ed. by M. Kaltenbach and H. N. Neufeld, pp. 17-25, Excerpta Medica, Amsterdam — Oxford — Princeton, 1982.

- 6) *Taira, N.*: Pharmacological profile of diltiazem, a calcium entry blocker. In: Calcium Modulators, ed. by T. Godfraind, A. Albertini and R. Paoletti, pp. 339–350, Elsevier Biomedical Press, Amsterdam – New York – Oxford, 1982.
- 7) Endoh, M., Maruyama, M. and *Taira, N.*: Adenosine-induced changes in rate of beating and cyclic nucleotide levels in rat atria: Modification by islet-activating protein. In: Physiology and Pharmacology of Adenosine Derivatives, ed. by J. W. Daly, Y. Kuroda, J. W. Phillis, H. Shimizu, and M. Ui, pp. 127–141, Raven Press, New York, 1983.
- 8) *Taira, N.*: Verapamil, diltiazem, nifedipine — Comparative studies on the AV node. In: Drug Development and Evaluation Vol. 9: New Calcium Antagonists: Recent Development and Prospects, ed. by A. Fleckenstein, K. Hashimoto, M. Herrmann, A. Schwartz and L. Seipel, pp. 37–52, Gustav Fischer Verlag, Stuttgart – New York, 1983.
- 9) *Taira, N.*: New positive inotropic agents: Their modes and mechanisms of action. In: Regulation of Cardiac Function, ed. by H. Abe, Y. Ito, M. Tada and L. H. Opie, pp. 293–303, Japan Sci. Soc. Press, Tokyo/VNU Sci. Press, Utrecht, 1984.
- 10) *Taira, N.*: Differences between dihydropyridine and non-dihydropyridine calcium antagonists in inotropic, chronotropic, dromotropic and vascular effects: Differences in underlying mechanism. In: Bayer Symposium K. Cardiovascular Effects of Dihydropyridine-Type Calcium Antagonists and Agonists, ed. by A. Fleckenstein, C. Van Breemen, R. Groß and F. Hoffmeister, pp. 370–378, Springer-Verlag, Berlin – Heidelberg – New York – Tokyo, 1985.
- 11) Endoh, M., Yanagisawa, T., *Taira, N.* and Blinks, J. R.: Effects of new inotropic agents on cyclic nucleotide metabolism and calcium transients in canine ventricular muscle. *Circulation* 73 (Suppl. III): 117–131, 1986.
- 12) *Taira, N.*: Differences in cardiovascular profile among calcium antagonists. *Am. J. Cardiol.* 59: 24B–29B, 1987.
- 13) *Taira, N.*: An overview of the pharmacology of procaterol. *J. Resp. Dis.* 8 (Suppl.): S22–S24, 1987.
- 14) *Taira, N.*: Similarity and dissimilarity in the mode and mechanism of action between nicorandil and classical nitrates: An overview. *J. Cardiovasc. Pharmacol.* 10 (Suppl. 8): S1–S9, 1987.

- 15) *Taira, N.*: Inotropic agents: New positive inotropic agents and their mechanism of action. In: The Proceeding of the 10th International Congress of Pharmacology, ed. by M. J. Rand and C. Raper, pp. 477-480, Elsevier Science Publishers, Amsterdam, 1987.
- 16) *Taira, N.*: Nicorandil as a hybrid between nitrates and potassium channel activators. *Am. J. Cardiol.* 63: 18J-24J, 1989.
- 17) Yanagisawa, T., Ishii, K., Yokoyama, H., Kurosawa, H. and *Taira, N.*: Selective stimulation of high-affinity β_1 -receptors: Its implication in the treatment of mild heart failure. In: Recent Progress in Failing Heart Syndrome, ed. by S. Sasayama and H. Suga, pp. 281-302, Springer-Verlag, Tokyo - Berlin - Heidelberg - New York - London - Paris - Hong Kong - Barcelona, 1991.
- 18) Yanagisawa, T. and *Taira, N.*: Coronary vasodilation by K^+ channel openers. In: Ion Channels of Vascular Smooth Muscle Cells and Endothelial Cells, ed. by N. Sperelakis and H. Kuriyama, pp. 151-162, Elsevier, New York - Amsterdam - London - Tokyo, 1991.
- 19) *Taira, N.*: Nifedipine preclinical and clinical overview. In: Nifedipine Quarter Century Symposium Tokyo '93, ed. by N. Taira, S. Hosoda, P. R. Lichtlen, pp. 3-9, Excerpta Medica, Amsterdam - Hong Kong - Princeton - Sydney - Tokyo, 1994.

(和 文)

- 1) 平 則夫. Ca-antagonists の薬理. Ca-antagonists — 薬理と臨床. 田辺製薬株式会社, 大阪, pp.1-35, 1979.
- 2) 平 則夫. 内臓における痛みの受容. 痛み — 基礎と臨床. 市岡正道, 中浜 博, 山村秀夫編集, 朝倉書店, 東京, pp.39-48, 1980.
- 3) 平 則夫. 発痛物質と秩序. ペインクリニック — 各種のいたみとその取り扱い方. 内科セミナー PN4. 織田敏次他 9 名編集, 永井書店, 大阪, pp.33-47, 1980.
- 4) 平 則夫. Ca^{++} 拮抗剤の薬理. Ca^{++} 拮抗剤の基礎と臨床. メディカルトリビューン日本支社, 東京, pp.35-52, 1981.
- 5) 平 則夫. Ca^{++} 拮抗薬のプロファイル. カルシウム拮抗薬 — 基礎と臨床. 木村栄一, 平 則夫編集, 医薬ジャーナル社, 大阪, pp.13-30, 1983.

- 6) 佐藤慶祐, 平 則夫. Ca^{++} 拮抗薬の冠循環に及ぼす影響. カルシウム拮抗薬 — 基礎と臨床.
木村栄一, 平 則夫編集, 医薬ジャーナル社, 大阪, pp.107-123, 1983.
- 7) 平 則夫. 冠循環の生理と薬理. 狭心症 — 冠動脈攣縮の基礎と臨床.
橋本虎六監修, 羊土社, 東京, pp.29-42, 1984.
- 8) 柳澤輝行, 平 則夫. カテコールアミン受容体サブタイプの機能的役割. 内因性神経活性物質.
吉田 博編著, 中外医学社, pp.176-186, 1989.
- 9) 平 則夫. 血管とカルシウム. 心臓と血管の科学. 第4回「大学と科学」
公開シンポジウム組織委員会編, クバプロ, 東京, pp.180-193, 1990.
- 10) 柳澤輝行, 平 則夫. 細胞内 Ca イオン濃度の測定. 血管研究法の進歩.
江橋節郎編集, 学会出版センター, pp.277-289, 1990.
- 11) 柳澤輝行, 勅使河原義, 森 豊樹, 平 則夫. エンドセリンの血管収縮作用に
対する血管拡張薬の影響. エンドセリン — 最新情報.
真崎知生監修, 平田結喜緒, 柳沢正史編集, 中外医学社, pp.88-92, 1990.
- 12) 柳澤輝行, 平 則夫. K^+ チャネルとモジュレーター. カルシウムと情報伝達系
— 平滑筋研究の現状と実験法.
浦川紀元, 唐木英明編集, 文永堂出版, 東京, pp.69-81, 1991.